Omica Labs

PANEL DOLOR

Listado de genes

ABCA1[1],ABCB1[1],ABCG1[1],ADCYAP1[1],ADORA1[1],ADORA2A[1],ADORA3[1],ADRA2A[1],ADRB2[1],AGTR2[1],ALOX12[1],AQP4[1],AR[1],ARNTL[1],ARRB2[1],ASIC1[1],ASIC3[1],ATAT1[1],ATF4[1],ATL1[2],ATL3[3],AVIL[1],BACE1[1],BAX[1],BDKRB1[1],BDKRB2[1],BDNF[1],BEGAIN[1],BHLHA9[1],CACNA1B[1],CACNA2D1[1],CACNA2D3[1],CACNA2D3[1],CACNA2D3[1],CACNB3[1],CACNG2[1],CALCA[1],CAMK2A[1],CAMKIV[1],CAPN1[1],CCKBR[1],CCR2[1],CDK5R1[1],CDKL5[1],CHRM2[1],CLCN6[1],CNR1[1],CNR2[1],COMT[1],COX1[1],COX2[1],CRIP2[1],CSF1[1],CSF2[1],CTSB[1],CTSS[1],CYP19A1[1],CYP2D6[4],CYP2D19[5],DAN[1],DAO[1],DBH[1],DDO[1],DICER1[1],DLG2[1],DMPIEZO[1],DPP4[1],EAR2[1],EDN1[1],EDNRB[1],EFNB1[1],EFNB2[1],EGR1[1],EHMT2[1],EIF2[1],EIF2AK2[1],EIF2AK3[1],EIF4E[1],EIF4EBP1[1],ESR1[1],F2RL1[1],FAAH[1],FKBP5[1],FPR1[1],GABBR1[1],GABRG2[1],GAL[1],GALR1[1],GALR2[1],GCH1[1],GCH1[1],GFRA2[1],GNA11[1],GNAO1[1],GNAQ[1],GNAZ[1],GPR34[1],GPR55[1],GRIK1[1],GRIN1[1],GRIN2A[1],GRIN2B[1],GRK2[1],GRK3[1],GRK5[1],GRM1[1],GRM5[1],HAMLET[1],HCN2[1],HDC[1],HMOX2[1],HOXB8[1],HRH1[1],HRH2[1],IAPP[1],ICA1[1],IFNGR[1],IL10[1],IL10R1[1],IL1A[1],IL1B[1],IL1R1[1],IL23[1],IL23R[1],IL4[1],IL6[1],IL6ST[1],KCNA2[1],KCNJ3[1],KCNJ6[1],KCNK2[1],KCNS1[1],KLF11[1],KIF1A[6],LEP[1],LMX1B[1],LPAR1[1],LPAR5[1],LRP1[1],MAPK1[1],MAPK10[1],MAPK3[1],MAPK8[1],MAPK9[1],MC1R[1],MDK[1],MECP2[1],MKK3[1],MKK6[1],MME[1],MMP13,MMP24[1],MMP9[1],MNK1[1],MNK2[1],MPZ[1],MRGPRE[1],MTA1[1],NDN[1],NF1[1],NGF[1],NGFR[1],NGR2[1],NLGN2[1],NOS1[1],NOS2[1],NPTX1[1],NPY[1],NPY1R[1],NRG1[1],NTRK1[1],NTSR1[1],NTSR2[1],OPRD1[1],OPRK1[1],OPRM1[1],OSM[1],P2RX3[1],P2RX7[1],P2RY12[1],PANX1[1],PDE10A[1],PDYN[1],PENK1[1],PER2[1],PIP5K1C[1],PIRT[1],PLCB3[1],PLP1[1],PNOC[1],POMC[1],PPARA[1],PPP1R14C[1],PRDM12[1],PRKAR1B[1],PRKCG[1],PRKG1[1],PRLR[1],PROK2[1],PROKR1[1],PRRX1[1],PRX[1],PTGER1[1],PTGER3[1],PTGIR[1],PTN[1],PTPN5[1],RAG1[1],REST[1],RGS9[1],RLUA-1[1],RLUA-2[1],RUNX1[1],S100A10[1],SARM1[1],SCH1[1],SCN10A[1],SCN11A[1],SCN3A[1],SCN9A[1],SHANK3[1],SLC12A2[1],SLC12A5[1],SLC17A6[1],SMAL[1],SORT1[1],SPARC[1],SPP1[1],ST8SIA1[1],SIRT1[7],TAC1[1],TAC4[1],TACR1[1],TBK1[1],TDAG8[1],TLR4[1],TMEM16F[1],TMEM35A[1],TNF[1],TNFRSF1A[1],TNFRSF1B[1],TRPA1[1],TRPC5[1],TRPM2[1],TRPM8[1],TRPV1[1],TRPV4[1],TSC2[1],VGLUT2[1].

Anexo I- Bibliografía

1. Wistrom, E., Chase, R., Smith, P. R. & Campbell, Z. T. A compendium of validated pain genes. *WIREs Mechanisms of Disease* **14**, e1570 (2022).
2. Guelly, C. *et al.* Targeted high-throughput sequencing identifies mutations in atlastin-1 as a cause of hereditary sensory neuropathy type I. *Am. J. Hum. Genet.* **88**, (2011).
3. Kornak, U. *et al.* Sensory neuropathy with bone destruction due to a mutation in the membrane-shaping atlastin GTPase 3. *Brain* **137**, (2014).
4. Jannetto, P. J. & Bratanow, N. C. Pharmacogenomic considerations in the opioid management of pain. *Genome Med.* **2**, 66 (2010).
5. Hicks, J. K. *et al.* Clinical pharmacogenetics implementation consortium guideline (CPIC) for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants: 2016 update. *Clin. Pharmacol. Ther.* **102**, 37–44 (2017).
6. Rivière, J. B. *et al.* KIF1A, an axonal transporter of synaptic vesicles, is mutated in hereditary sensory and autonomic neuropathy type 2. *Am. J. Hum. Genet.* **89**, (2011).
7. Wang, Y. *et al.* SIRT1 mediates the excitability of spinal CaMKIIα-positive neurons and participates in neuropathic pain by controlling Nav1.3. *CNS Neurosci. Ther.* **30**, (2024).